



ENVIRONMENTAL  
HEALTH  
PERSPECTIVES

<http://www.ehponline.org>

## Birth Outcomes and Maternal Residential Proximity to Natural Gas Development in Rural Colorado

**Lisa M. McKenzie, Ruixin Guo, Roxana Z. Witter,  
David A. Savitz, Lee S. Newman, and John L. Adgate**

<http://dx.doi.org/10.1289/ehp.1306722>

**Received: 27 February 2013**

**Accepted: 28 January 2014**

**Advance Publication: 28 January 2014**

# **Birth Outcomes and Maternal Residential Proximity to Natural Gas Development in Rural Colorado**

Lisa M. McKenzie,<sup>1</sup> Ruixin Guo,<sup>2</sup> Roxana Z. Witter,<sup>1</sup> David A. Savitz,<sup>3</sup> Lee S. Newman,<sup>1</sup> and John L. Adgate<sup>1</sup>

<sup>1</sup>Department of Environmental and Occupational Health, Colorado School of Public Health, Aurora, Colorado, USA; <sup>2</sup>Department of Biostatistics and Informatics, Colorado School of Public Health, Aurora, Colorado, USA; <sup>3</sup>Department of Epidemiology, Brown University, Providence, Rhode Island, USA

**Address correspondence to** Lisa McKenzie, Department of Environmental and Occupational Health, Colorado School of Public Health, 13001 E. 17th Place, Campus Box B119, Aurora, Colorado 80045 USA. Telephone: 303.724.5557. E-mail: Lisa.McKenzie@ucdenver.edu

**Running Title:** Natural gas development and birth outcomes

**Acknowledgments:** This study was supported by the Department of Environmental and Occupational Health at the Colorado School of Public Health. The Colorado Department of Public Health (CDPHE) and Environment's Health Statistics and Colorado Responds to Children with Special Needs Sections provided outcome data for this study. CDPHE specifically disclaims responsibility for any analyses, interpretations, or conclusions.

**Competing Financial Interests:** The authors declare they have no competing financial interests.

## **Abstract**

**Background:** Birth defects are a leading cause of neonatal mortality. Natural gas development (NGD) emits several potential teratogens and US production is expanding.

**Objectives:** We examined associations between maternal residential proximity to NGD and birth outcomes in a retrospective cohort study of 124,842 births between 1996 and 2009 in rural Colorado.

**Methods:** We calculated inverse distance weighted natural gas well counts within a 10-mile radius of maternal residence to estimate maternal exposure to NGD. Logistic regression, adjusted for maternal and infant covariates, was used to estimate associations with exposure tertiles for congenital heart defects (CHDs), neural tube defects (NTDs), oral clefts, preterm birth, and term low birth weight. The Association with term birth weight was investigated using multiple linear regression.

**Results:** Prevalence of CHDs increased with exposure tertile, with an odds ratio (OR) of 1.3 for the highest tertile (95% CI: 1.2, 1.5) and NTD prevalence was associated with the highest tertile of exposure (OR = 2.0, 95% CI: 1.0, 3.9, based on 59 cases), compared to no gas wells within a 10-mile radius. Exposure was negatively associated with preterm birth and positively associated with fetal growth, though the magnitude of association was small. No association was found between exposure and oral clefts.

**Conclusions:** In this large cohort, we observed an association between density and proximity of natural gas wells within a 10-mile radius of maternal residence and prevalence of CHDs and possibly NTDs. Greater specificity in exposure estimates are needed to further explore these associations.

## Introduction

Approximately 3.3% of US live births have a major birth defect (CDPHE 2011; Parker et al. 2010) which account for 20% of infant deaths as well as 2.3% of premature death and disability (McKenna et al. 2005). Oral clefts, neural tube defects (NTDs), and congenital heart defects (CHD) are the most common classes of birth defects (Parker et al. 2010). These defects are thought to originate in the first trimester as a result of polygenic inherited disease or gene-environment interactions (Brent 2004). Suspected non-genetic risk factors for these birth defects include folate deficiency (Wald and Sneddon 1991), maternal smoking (Honein et al. 2006), alcohol abuse and solvent use (Romitti et al. 2007), and exposure to benzene (Lupo et al. 2010a; Wennborg et al. 2005), toluene (Bowen et al. 2009), polycyclic aromatic hydrocarbons (PAHs) (Ren et al. 2011), and petroleum based solvents, including aromatic hydrocarbons (Chevrier et al. 1996). Associations between air pollution (Volatile Organic Compounds [VOCs], particulate matter [PM] and nitrogen dioxide [NO<sub>2</sub>]) and low birth weight and preterm birth have been reported (Ballester et al. 2010; Brauer et al. 2008; Dadvand et al. 2013; Ghosh et al. 2012; Llop et al. 2010). Many of these air pollutants are emitted during development and production of natural gas (referred to herein as NGD) and concerns have been raised that they may increase risk of adverse birth outcomes and other health effects (Colborn et al. 2011; McKenzie et al. 2012). Increased prevalence of low birth weight and small for gestational age, as well as reduced APGAR scores were reported in infants born to mothers living near NGD in Pennsylvania (Hill 2012).

Technological advances in directional drilling and hydraulic fracturing have resulted in a global boom of drilling and production of natural gas reserves (U. S. EIA 2011a, b; Vidas and Hugman

2008). NGD is an industrial process resulting in potential worker and community exposure to multiple environmental stressors (Esswein et al. 2013; King 2012; Witter et al. 2013). Diesel powered heavy equipment is used for worksite development as well as transporting large volumes of water, sand, and chemicals to sites and for waste removal (Witter et al 2013). It is increasingly common for NGD to encroach on populated areas, potentially exposing more people to air and water emissions as well as noise and community level changes that may arise from industrialization (COGCC 2009). Studies in Colorado, Texas, Wyoming, and Oklahoma have demonstrated that NGD results in emission of VOCs, NO<sub>2</sub>, SO<sub>2</sub>, PM, and PAHs from either the well itself or from associated drilling processes or related infrastructure, i.e., drilling muds, hydraulic fracturing fluids, tanks containing waste water and liquid hydrocarbons, diesel engines, compressor stations, dehydrators, and pipelines (CDPHE 2007; Frazier 2009; Kemball-Cook et al. 2010; Olaguer 2012; Walther 2011; Zielinska et al. 2011). Some of these pollutants, e.g., toluene, xylenes, and benzene, are suspected teratogens (Lupo et al. 2010; Shepard 1995) or mutagens (ATSDR 2007) and are known to cross the placenta (Bukowski 2001), raising the possibility of fetal exposure to these and other pollutants resulting from NGD. Currently, there are few studies on the effects of air pollution or NGD on birth outcomes.

This analysis explores the association between maternal exposure to NGD and birth outcomes, using a dataset with individual-level birth data and geocoded natural gas well locations. We conducted a retrospective cohort study to investigate the association between density and proximity of natural gas wells within a 10-mile radius of maternal residences in rural Colorado and three classes of birth defects, preterm birth, and fetal growth.

## **Methods**

### **Study population**

We used information available in the publically accessible Colorado Oil and Gas Information System to build a geocoded dataset with latitude, longitude, and year of development (1996-2009) for all gas wells in rural Colorado (COGIS 2011). Live birth data were obtained from the Colorado Vital Birth Statistics (CDPHE 2011). Geocoded maternal addresses at time of birth were linked to the well locations. Distance of each maternal residence from all existing (not abandoned) natural gas wells within a 10-mile radius was then computed using spherically-adjusted straight line distances. We conducted our analysis on the final de-identified database containing maternal and birth outcome data described below and distance to all wells within the 10- mile radius. The Colorado Multiple Institutional Review Board reviewed and approved our study protocol. Informed consent was not required.

We restricted analysis to births occurring from 1996 to 2009 to focus our analysis on growth of unconventional NGD, characterized by use of hydraulic fracturing and/or directional drilling (King 2012), which expanded rapidly in Colorado beginning around 2000 (COGIS 2011). We also restricted our analysis to rural areas and towns with populations of less than 50,000 (excluding the Denver-Metropolitan area, El Paso County and the cities of Fort Collins, Boulder, Pueblo, Grand Junction and Greeley) in 57 counties to reduce potential for exposure to other pollution sources, such as traffic, congestion, and industry. The final study area included locations with and without NGD. We conducted a retrospective study on the resulting cohort of 124,842 live births to explore associations between birth outcomes and exposure to NGD

operations. We restricted eligibility to singleton births and excluded the small proportion (< 5%) of non-white births because there were too few to analyze separately.

### **Birth outcomes**

Identified birth outcomes were (1) oral cleft, including cleft lip with and without cleft palate as well as cleft palate (ICD9-CM 749.xx) (National Center for Health Statistics 2011); (2) NTD, including anencephalus, spina bifida without anecephaly, and encephalocele (ICD9-CM740.xx, ICD9-CM741.xx, and ICD9-CM742.0); (3) CHD, including transposition of great vessels, tetralogy of Fallot, ventricular septal defect, endocardial cushion defect, pulmonary valve atresia and stenosis, tricuspid valve atresia and stenosis, Ebstein's anomaly, aortic valve stenosis, hypoplastic left heart syndrome, patent ductus arteriosus, coarctation of aorta, and pulmonary artery anomalies (ICD9-CM745.xx, ICD9-CM-746.xx, ICD9-CM747.xx, excluding 746.9, 747.5); (4) preterm birth (< 37 weeks completed gestation); (5) term low birth weight ( $\geq$  37 weeks completed gestation and birth weight < 2500 grams); and (6) term birth weight as a continuous measure. Births with an oral cleft, NTD, or CHD were excluded from preterm birth and term low birth weight analysis. Preterm births were excluded from term birth weight analysis. Oral cleft, CHD, and NTD cases in the Colorado Responds to Children with Special Needs (CRCSN) birth registry, obtained from hospital records, the Newborn Genetics Screening Program, the Newborn Hearing Screening Program, laboratories, physicians, and genetic, developmental and other specialty clinics (CRCSN 2011), were matched with Colorado live birth certificates. Cases are reflective of reporting as of July 12, 2012, were not necessarily confirmed by medical record review, and are subject to change as CRCSN ascertains diagnosis up to 3 years old and/or supplements information by medical record review. We analyzed birth defects in three heterogeneous groups to increase statistical power. Dataset information was not

sufficient to distinguish between multiple and isolated birth anomalies or to identify chromosomal birth anomalies. In an exploratory analysis, we considered seven clinical diagnostic groupings of CHDs: (1) conotruncal defects (Tetralogy of Fallot and transposition of great vessels); (2) endocardial cushion and mitrovalve defects (EMD: endocardial cushion defect and hypoplastic left heart syndrome); (3) pulmonary artery and valve defects (PAV: pulmonary valve atresia and stenosis and pulmonary artery anomalies); (4) tricuspid valve defects (TVD: tricuspid valve atresia and stenosis and Ebstein's anomaly); (5) aortic artery and valve defects (aortic valve stenosis and coarctation of aorta); (6) ventricular septal defects (VSD); and (7) patent ductus arteriosus in births greater than 2500 grams (Gilboa 2005).

### **Exposure assessment**

Distribution of the wells within a 10-mile radius of maternal residence shows 50% and 90% of wells to be within 2.3 and 7.7 miles of maternal residence, respectively. We used an inverse distance weighted (IDW) approach, commonly used to estimate individual air pollutant exposures from multiple fixed locations (Brauer 1998, Ghosh 2012), to estimate maternal exposure. Our IDW well count accounts for the number of wells within the 10-mile radius of the maternal residence, as well as distance of each well from the maternal residence, giving greater weight to wells closest to the maternal residence. For example, an IDW well count of 125 wells per mile could be computed from 125 wells each located 1 mile from the maternal residence or 25 wells each located 0.2 miles from the maternal residence. We calculated the IDW well count of all existing natural gas wells in the birth year within a 10-mile radius of each maternal residence as a continuous exposure metric:

$$IDW \text{ well count} = \sum_{i=1}^n \frac{1}{a_i} \quad [1]$$



where

IDW well count = IDW count of existing wells within a 10-mile radius of maternal residence in the birth year

$d_i$  = distance of the  $i^{\text{th}}$  individual well from maternal residence

$n$  = number of existing wells within a 10-mile radius of maternal residence in the birth year.

The IDW well count was calculated for each maternal residence with  $\geq 1$  gas wells within 10 miles. The final distribution then was divided into tertiles (low, medium, and high) for subsequent logistic and linear regression analysis. Each tertile was compared to the referent group (no natural gas wells within 10 miles, IDW well count = 0).

### **Statistical analysis**

We used logistic regressions to study associations between each dichotomous outcome and IDW exposure group. We also considered term birth weight as a continuous outcome using multiple linear regression. First, we estimated the crude odds ratio (OR) associated with IDW exposure tertile for each binary outcome, followed by a Cochran-Armitage test to evaluate linear trends in binominal proportions with increasing IDW exposure (none, low, medium, and high). We further investigated associations by adjusting for potential confounders, as well as infant and maternal covariates selected based on both *a priori* knowledge and empirical consideration of their association with exposure and an outcome. Specifically, covariates in our analysis of all outcomes except outcomes with very few events (i.e., NTDs, CTDs, EMDs, and TVDs) included maternal age, education (< 12, 12, 13-15,  $\geq 16$  years), tobacco use (smoker, non-smoker), ethnicity (Hispanic, non-Hispanic), and alcohol use (yes, no), as well as parity at time of

pregnancy (0, 1, 2, > 2) and infant gender. Gestational age was also included in the analysis of term birth weight. Elevation of maternal residence also was considered in the analysis because most wells are below 7000 feet, and elevation has been associated with both preterm birth and low birth weight (Niermeyer et al. 2009). For 272 births where elevation of maternal residence was missing, elevation was imputed using mean elevation for maternal zip code. For outcomes with very few events, only elevation was included in the multiple logistic modeling to avoid unstable estimates. The ORs and their 95% confidence intervals (CI) were used to approximate relative risks for each outcome associated with IDW count exposure tertile (low, medium, and high) compared to no wells within 10 miles, which is reasonable because of the rarity of the outcomes. We considered the statistical significance of the association, as well as the trend, in evaluating results, at an alpha of 0.05. We evaluated the confounding potential of the 1998 introduction of folic acid fortification on the birth defect outcomes and found only a decrease in NTD prevalence after 1998 (Supplemental Material, Table S1).

In a sensitivity analyses, we explored reducing exposure to 2- and 5-mile buffers around the maternal residence, as well as restricting the cohort to births occurring between 2000 and 2009 to exclude births before the expansion of NGD. We report estimated associations with 95% CIs. All statistical analyses were conducted using SAS® software version 9.3 (Cary, NC).

## **Results**

Births were approximately evenly divided between exposed and unexposed groups (0 wells in a 10-mile radius versus  $\geq 1$  well in a 10-mile radius) (Table 1). Estimated exposure, as represented by IDW well counts, tended to be higher for births to mothers with residence addresses at lower elevations (< 6000 feet), and among nonsmoking and Hispanic mothers (Table 1).

Both crude and adjusted estimates indicate a monotonic increase in the prevalence of CHDs with increasing exposure to NGD, as represented by IDW well counts (Table 2). Births to mothers in the most exposed tertile (> 125 wells/mile) had a 30% greater prevalence of CHDs (95% CI: 1.2, 1.5) than births to mothers with no wells within a 10-mile radius of their residence.

Prevalence of NTDs was positively associated with only the third exposure tertile, based on crude and estimated adjusted ORs for elevation (Table 2). Births in the highest tertile (> 125 well/mile) were 2.0 (95% CI: 1.0, 3.9) times more likely to have a NTD than those with no wells within a 10-mile radius, based on 59 available cases. We observed no statistically significant associations between oral clefts and NGD, based on trend analysis across categorical IDW well count exposure (Table 2).

Both crude and adjusted estimates for preterm birth suggest a slight (< 10%) decreased risk of preterm birth with increasing exposure to NGD (Table 3). Crude term low birth weight measures suggested decreased risk of term low birth weight with increasing exposure to NGD. A weak non-linear trend remained after adjusting for elevation and other covariates. This association is consistent with the multiple linear regression results for continuous term birth weight, in which mean birth weights were 5-24 grams greater in the higher IDW well count exposure tertiles than the referent group.

We observed a monotonic increase in the prevalence of NTDs with increasing exposure to NGD in our sensitivity analyses using 2- and 5-mile exposure radii as well as some attenuation in decreased risk for preterm birth and term low birth weight (Supplemental Material, Tables S2-7). Restricting births to 2000 to 2009, the period of most intense NGD in Colorado, attenuated the

positive association between NTDs in the highest tertile and did not alter observed relationships for other birth outcomes (Supplemental Material, Tables S2-S7).

Exploratory analysis of CHDs by clinical diagnostic groups indicates increased prevalence of PAV defects by 60% (95% CI 1.1, 2.2), VSDs by 50% (95% CI 1.1, 2.1), and TVDs by 400% (95% CI 1.3, 13) in the most exposed tertile compared to those with no wells within a 10-mile radius (Table 4).

## **Discussion**

We found positive associations between density and proximity of natural gas wells within a 10-mile radius of maternal residence and birth prevalence of CHDs and possibly NTDs. Prevalence of CHDs increased monotonically from the lowest to highest exposure tertile, though even in the highest tertile, the magnitude of the association was modest. Prevalence of NTDs was only elevated in the highest tertile of exposure. We also observed small negative associations between density and proximity of natural gas wells within a 10-mile radius of maternal residence and preterm birth and term low birth weight, and a small positive association with mean birth weight. We found no indication of an association between density and proximity of natural gas wells within a 10-mile radius of maternal residence and oral cleft prevalence.

Non-genetic risk factors for CHDs and NTDs possibly attributable to NGD include maternal exposure to benzene (Lupo et al. 2010; Wennborg et al. 2005), PAHs (Ren et al. 2011), solvents (Brender et al. 2002; Chevrier et al. 1996; Desrosiers et al. 2012; McMartin et al. 1998), and air pollutants (NO<sub>2</sub>, SO<sub>2</sub>, PM) (Vrijheid 2011). NGD emits multiple air pollutants, including benzene and toluene, during the “well completion” phase (when gas and water flow back to the

surface after hydraulic fracturing) as well as from related infrastructure (CDPHE 2009a, b; GCPH 2009; Gilman et al. 2013; McKenzie et al. 2012; Pétron et al. 2012). Ambient benzene levels in areas with active NGD in Northeast Colorado ranged from 0.03 to 6 parts per billion by volume (ppbv) (CDPHE 2012; Gilman et al. 2013; Pétron et al. 2012). Furthermore, 24-hour average ambient air benzene levels near active well development sites in Western Colorado ranged from 0.03 to 22 ppbv (McKenzie et al. 2012).

Two previous case-control studies have reported associations between maternal exposure to benzene and birth prevalence of NTDs and/or CHDs (Lupo et al. 2010a; Wennborg et al. 2005). Lupo's study of 4531 births in Texas found that mothers living in census tracts with the highest ambient benzene levels (0.9 – 2.33 ppbv) were 2.3 times more likely to have offspring with spina bifida than mothers living in census tracts with the lowest ambient benzene levels (95% CI: 1.22, 4.33) (Lupo et al. 2010a). An occupational study of Swedish laboratory employees found a significant association between exposure to occupational levels of benzene in the critical window between conception, organogenesis, and neural crest formation and neural crest malformations (Wennborg et al. 2005). Children born to 298 mothers exposed to benzene had 5.3 times greater prevalence of neural crest malformations than children born to mothers not exposed to benzene (95% CI: 1.4, 21.1). Other studies of maternal exposures to organic solvents, some of which contain benzene, have reported associations between maternal occupational exposure to organic solvents and major birth defects (Brender et al. 2002; McMartin et al. 1998; Desrosiers et al. 2012). While exposure to benzene is a plausible explanation for the observed associations, further research is needed to examine whether these associations are replicated and whether benzene specifically explains these associations.

Air pollutants emitted from diesel engines used extensively in NGD also may be associated with CHDs and/or NTDs. Trucks with diesel engines are used to transport supplies, water, and waste to and from gas wells, with 40 to 280 truck trips per day per well pad during development (Witter et al. 2013). Generators equipped with diesel engines are used both to drill wells and in hydraulic fracturing. Air pollutants in diesel exhaust include NO<sub>2</sub>, SO<sub>2</sub>, PM, and PAHs. A meta-analysis of four studies suggested associations of maternal NO<sub>2</sub> and SO<sub>2</sub> exposures with coarctation of the aorta and tetralogy of Fallot, and of maternal PM<sub>10</sub> exposure with arterial septal defects (Vrijheid et al. 2011). Two case-control studies in China reported positive associations between PAH concentrations in maternal blood and the placenta and NTDs (Li et al. 2011; Naufal et al. 2010). Several CHDs were associated with traffic related carbon monoxide and ozone pollution in a case control study of births from 1987 to 1993 in Southern California (Ritz et al. 2002).

The small negative associations with term low birth weight and preterm birth in our study population were unexpected given that other studies have reported positive associations between these outcomes and urban air pollution (Ballester et al. 2010; Brauer et al. 2008; Dadvand et al. 2013; Ghosh et al. 2012; Llop et al. 2010) and proximity to natural gas wells (Hill 2012). It is possible that rural air quality near natural gas wells in Colorado is not as compromised as urban air quality in these studies and exposure represented as IDW well count may not well represent air quality. In addition, the power of our large cohort increases the likelihood of false positive results for small associations close to the null. While associations were consistent across measures of birth weight, i.e., reduced risk of term low birth weight and increase in mean birth weight, they attenuated towards the null in sensitivity analysis for 2 and 5 mile radii

(Supplemental Tables S6-S7). If causal, stronger associations would be expected with more stringent exposure definitions. Our incomplete ability to adjust for socioeconomic status, health, nutrition, prenatal care, and pregnancy complications likely accounts for these unexpected findings.

This study has several limitations inherent in the nature of the available data. Not all birth defects were confirmed by medical record review. Also, birth defects are most likely undercounted, because non-live births, terminated pregnancies, and later life diagnoses (after age 3 years) are not included. Birth weight and gestational age were obtained from birth certificates, which are generally accurate for birth weight and useful but less accurate for gestational age (DiGiuseppe et al. 2002). Data on covariates were obtained from birth certificates and were limited to basic demographic, education, and behavioral information available in the vital records. Distribution of covariates among exposure tertiles and the unexposed group was similar; nevertheless, our incomplete ability to adjust for socioeconomic status, health, nutrition, prenatal care, and pregnancy complications may have resulted in residual confounding. In addition, low event outcomes (e.g., NTDs) were only adjusted for elevation. The dataset did not contain information on maternal folate consumption and genetic anomalies, both independent predictors of our outcomes, which may have confounded these results. We did observe a large decrease in the prevalence of NTDs after the introduction of folic acid in 1998 and small increases in the prevalence of CHDs, and oral clefts albeit none of the estimates are statistically significant (supplemental table S1). Further study is needed to determine if unaccounted folate confounding is attenuating our results towards the null. There is no evidence indicating genetic anomalies would differ by IDW well count around maternal residence.

Due to rarity of specific birth defects in the study population, birth defects were aggregated into three general groups. This limited our study in that associations with specific birth defects, if present, may have been obscured. An exploratory analysis of CHDs by clinical diagnostic groups indicates increased prevalence of specific diagnostic groups (i.e., PAV, VSD, and TSV) compared to aggregated CHDs (Table 4).

Another limitation of this study is the lack of temporal and spatial specificity of the exposure assessment. Because we did not have maternal residential history, we assumed that maternal address at time of delivery was the same as maternal address during the first trimester of pregnancy, the critical time period for formation of birth defects. Studies in Georgia and Texas estimate that 22 to 30 percent of mothers move during their pregnancy and most mothers move within their locality (Miller et al. 2010 and Lupo et al. 2010b), potentially introducing some exposure misclassification for the early pregnancy period of interest. However, these studies found little difference in mobility between cases and controls (Miller et al. 2010, Lupo et al. 2010b) and maternal mobility did not significantly influence the assessment of benzene exposure (Lupo et al. 2010b). We were only able to determine if a well existed within the calendar year of birth (e.g., 2003) and did not have sufficient data to determine if a well existed within the first trimester of the pregnancy. Therefore, some non-differential exposure misclassification is likely and the overall effect of this is unknown.

Similarly, we only had consistent information on existence of a well in the birth year. Lack of information on natural gas well activity levels, such as whether or not wells were producing or undergoing development may have resulted in exposure misclassification. Actual exposure to natural gas-related pollutants likely varies by intensity of development activities. Lack of



temporal and spatial specificity of the exposure assessment would most likely have tended to weaken associations (Ritz et al. 2007; Ritz and Wilhelm 2008). To address spatial and temporal variability, additional air pollution measurements and modeling will be needed to improve exposure estimates at specific locations. Lastly, lack of information on the mother's activities away from her residence, such as work and recreation, as well as proximity of these activities to NGD were not available and may have led to further exposure misclassification and residual confounding.

## **Conclusion**

This study suggests a positive association between greater density and proximity of natural gas wells within a 10-mile radius of maternal residence and greater prevalence of CHDs and possibly NTDs, but not oral clefts, preterm birth, or reduced fetal growth. Further studies incorporating information on specific activities and production levels near homes over the course of pregnancy would improve exposure assessments and provide more refined effect estimates. Recent data indicate that exposure to NGD activities is increasingly common. The Colorado Oil and Gas Conservation Commission estimates that 26% of the more than 47,000 oil and gas wells in Colorado are located within 150 to 1000 feet of a home or other type of building intended for human occupancy (COGCC 2012). Taken together, our results and current trends in NGD underscore the importance of conducting more comprehensive and rigorous research on the potential health effects of NGD.

## References

- ATSDR (Agency for Toxic Substances and Disease Registry). 2007. Toxicological profile for benzene. <http://www.atsdr.cdc.gov/ToxProfiles/TP.asp?id=40&tid=14> [accessed 22 May 2013].
- Ballester F, Estarlich M, Iniguez C, Llop S, Ramon R, Esplugues A, et al. 2010. Air pollution exposure during pregnancy and reduced birth size: A prospective birth cohort study in Valencia, Spain. *Environ Health* 9:6.
- Bowen S, Irtenkauf S, Hannigna J, Stefanski A. 2009. Alterations in rat morphology following abuse patterns of toluene exposure. *Reprod Toxicol* 27:161-169.
- Brauer M, Lencar C, Tamburic L, Koehoorn M, Demers P, Karr C. 2008. A cohort study of traffic-related air pollution impacts on birth outcomes. *Environ Sci Technol* 116:680-686.
- Brender JD, Suarez L, Hendricks KA, Baetz RA, Larsen R. 2002. Parental occupation and neural tube defect-affected pregnancies among Mexican Americans. *J Occup Environ Med* 44:650-656.
- Brent RL. 2004. Environmental causes of human congenital malformations: The pediatrician's role in dealing with these complex clinical problems caused by a multiplicity of environmental and genetic factors. *Pediatrics* 113:957-968.
- Bukowski JA. 2001. Review of the epidemiological evidence relating toluene to reproductive outcomes. *Regul Toxicol and Pharmacol* 33:147-156.
- Chevrier C, Dananche B, Bahuaud M, Nelva A, Herman C, Francannet C, et al. 1996. Occupational exposure to organic solvent mixtures during pregnancy and the risk of non-syndromic oral clefts. *Occup Environ Med* 63:617-623.
- Colborn T, Kwiatkowski C, Schultz K, Bachran M. 2011. Natural gas operations from a public health perspective. *Hum Ecol Risk Assess* 17:1039-1056.
- CDPHE (Colorado Department of Public Health and Environment). 2007. Garfield County air toxics inhalation: Screening level human health risk assessment: Inhalation of volatile organic compounds measured in rural, urban, and oil & gas areas in air monitoring study (June 2005 – May 2007). <http://www.garfield-county.com/public-health/documents/Working%20Draft%20CDPHE%20Screening%20Level%20Risk%20Air%20Toxics%20Assessment%2012%2020%202007.pdf>. [accessed 22 May 2013].

- CDPHE. 2009a. Garfield County emissions inventory. [http://www.garfield-county.com/air-quality/documents/airquality/Garfield\\_County\\_Emissions\\_Inventory-2009.pdf](http://www.garfield-county.com/air-quality/documents/airquality/Garfield_County_Emissions_Inventory-2009.pdf) [accessed 22 May 2013].
- CDPHE 2009b. State of Colorado technical support document for recommended 8-hour ozone designations.  
<http://www.colorado.gov/cs/Satellite?blobcol=urldata&blobheadname1=Content-Disposition&blobheadname2=Content-Type&blobheadvalue1=inline%3B+filename%3D%22Recommended+8+Hour+Ozone+Area+Designations.pdf%22&blobheadvalue2=application%2Fpdf&blobkey=id&blobtable=MungoBlobs&blobwhere=1251808873080&ssbinary=true>. [accessed 22 May 2013].
- CDPHE. 2013. Colorado health information dataset birth statistics. Health Statistics Section.
- CDPHE. 2012. Air emissions case study related to oil and gas development in Erie, Colorado. Retrieved from: [http://www.colorado.gov/airquality/tech\\_doc\\_repository.aspx](http://www.colorado.gov/airquality/tech_doc_repository.aspx). [accessed 22 May 2013].
- COGCC (Colorado Oil and Gas Conservation Commission). 2009. Statement of basis, specific statutory authority, and purpose: New rules and amendments to current rules of the Colorado Oil and Gas Conservation Commission, 2 ccr 404-1. <http://cogcc.state.co.us/>. [accessed 22 May 2013].
- COGCC. 2012. Staff report. Colorado Department of Natural Resources. <http://cogcc.state.co.us/>. [accessed 22 May 2013].
- COGIS (Colorado Oil and Gas Information System) . 2011. Well production database. Vol. 2011. <http://cogcc.state.co.us/> [accessed 22 May 2013].
- CRCSN (Colorado Responds to Children with Special Needs). 2011. Birth defects dataset details. Available: <http://www.cdphe.state.co.us/cohid/crcsndata.html> [accessed October 6, 2011].
- Dadvand P, Parker JD, Bell ML, Bonzini M, Brauer M, Darrow L, et al. 2013. Maternal exposure to particulate air pollution and term birth weight: A multi-country evaluation of effect and heterogeneity. *Environ Health Perspect* 121:367-373.

- Desrosiers TA, Lawson CC, Meyer RE, Richardson DB, Daniels JL, Waters MA, et al. 2012. Maternal occupational exposure to organic solvents during early pregnancy and risks of neural tube defects and orofacial clefts. *Occup Environ Med* 69 493-499.
- DiGiuseppe DL, Aron DC, Ranbom L, Harper DL, Rosenthal GE. 2002. Reliability of birth certificate data: A multi-hospital comparison to medical records information. *Matern Child Health J* 6:169-179.
- Esswein EJ, Breitenstein M, Snawder J, Kiefer M, Sieber K. 2013. Occupational exposure to respirable crystalline silica during hydraulic fracturing. *J Occup Environ Hyg*, DOI:10.1080/15459624.2103.788352.
- Frazier A. 2009. Analysis of data obtained for the Garfield County air toxics study summer 2008. [http://www.garfield-county.com/air-quality/documents/airquality/2008\\_Targeted\\_Oil\\_and\\_Gas\\_Monitoring\\_Report.pdf](http://www.garfield-county.com/air-quality/documents/airquality/2008_Targeted_Oil_and_Gas_Monitoring_Report.pdf). [accessed 22 May 2013].
- GCPH (Garfield County Public Health Department). 2009. Garfield County 2009 air quality monitoring summary report. [http://www.garfield-county.com/air-quality/documents/airquality/2009\\_Air\\_Monitoring\\_Report.pdf](http://www.garfield-county.com/air-quality/documents/airquality/2009_Air_Monitoring_Report.pdf). [accessed 22 May 2013].
- Ghosh JKC, Wilhelm M, Su J, Goldberg D, Cockburn M, Jerrett M, et al. 2012. Assessing the influence of traffic-related air pollution on risk of term low birth weight on the basis of land-use-based regression models and measures of air toxics. *Am J Epidemiol* 175:1262-1274.
- Gilboa SM, Mendola P, Olshan AF, Langlois PH, Savitz DA, Loomis D, et al. 2005. Relation between ambient air quality and selected birth defects, Seven County Study, Texas, 1997-2000. *Am J Epidemiol* 162 (3):238-252.
- Gilman JB, Lerner BM, Kuster WC, de Gouw J. 2013. Source signature of volatile organic compounds (VOCs) from oil and natural gas operations in northeastern colorado. *Environ Sci Technol* 47 (3):1297-1305.
- Hill EL, 2012. Unconventional natural gas development and infant health: Evidence from Pennsylvania. Charles Dyson School of Applied Economics and Management at Cornell University. Work in Progress. <http://dyson.cornell.edu/research/researchpdf/wp/2012/Cornell-Dyson-wp1212.pdf>. [accessed 22 May 2013].

- Honein M, Rasmussen S, Reefhuis J, Romitti P, Lammer E, Sun L, et al. 2006. Maternal smoking and environmental tobacco smoke exposure and the risk of orofacial clefts. *Epidemiology* 18:226-233.
- Kemball-Cook S, Bar-Ilan A, Grant J, Parker L, Jung J, Santamaria W, et al. 2010. Ozone impacts of natural gas development in the Haynesville shale. *Environ Sci Technol* 44:9357-9363.
- Kim S, Vermeulen R, Waidyanatha S, Johnson BA, Lan Q, Smith MT, et al. 2006. Modeling human metabolism of benzene following occupational and environmental exposures. *Cancer Epidemiol Biomarkers Prev* 15:2246-2252.
- King GE. 2012. Hydraulic fracturing 101: what every representative, environmentalist, regulator, reporter, investor, university researcher, neighbor and engineer should know about estimating frac risk and improving frac performance in unconventional gas and oil wells. SPE 152596 presented at SPE Hydraulic Fracturing Technology Conference in The Woodlands, Texas, USA.
- Li Z, Zhang L, Ye R, Pei L, Liu J, Zheng X, et al. 2011. Indoor air pollution from coal combustion and the risk of neural tube defects in a rural population in Shanxi Province, China. *Am J Epidemiol* 174:451-458.
- Llop S, Ballester F, Estarlich M, Esplugues A, Rebagliato M, Iñiguez C. 2010. Preterm birth and exposure to air pollutants during pregnancy. *Environ Res* 110:778-785.
- Lupo P, Symanski E, Waller D, Chan W, Langlosi P, Canfield M, et al. 2010a. Maternal exposure to ambient levels of benzene and neural tube defects among offspring, Texas 1999-2004. *Environ Health Perspect* 119 (3): 397-402.
- Lupo PJ, Symanski E, Chan W, Mitchell LE, Waller DK, Canfield MA, et al. 2010b. Differences in exposure assignment between conception and delivery: the impact of maternal mobility. *Paediatr Perinat Epidemiol* 24: 200-208.
- McKenna MT, Michaud CM, Murray CJL, Marks JS. 2005. Assessing the burden of disease in the United States using disability-adjusted life years. *Am J Prev Med* 28:415-423.
- McKenzie LM, Witter RZ, Newman LS, Adgate JL. 2012. Human health risk assessment of air emissions from development of unconventional natural gas resources. *Sci Total Environ* 424:79-87.

- McMartin KI, Chu M, Kopecky E, Einarson TR, Koren G. 1998. Pregnancy outcome following maternal organic solvent exposure: A meta-analysis of epidemiologic studies. *Amn J Ind Med* 34:288-292.
- Miller A, Siffel Csaba, Correa A. 2010. Residential mobility during pregnancy: patterns and correlates. *Matern Child Health J* 14:625-634.
- National Center for Health Statistics. 2011. Classification of Diseases and Injuries. Retrieved from [ftp://ftp.cdc.gov/pub/Health\\_Statistics/NCHS/Publications/ICD-9/ucod.txt](ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Publications/ICD-9/ucod.txt). [accessed 22 May 2013].
- Naufal Z, Zhiwen L, Zhu L, Zhou GOD, McDonald T, He LY, et al. 2010. Biomarkers of exposure to combustion by-products in a human population in Shanxi, China. *J Expos Sci Environ Epidemiol* 20:310-319.
- Niermeyer S, Andrade Mollinedo P, Huicho L. 2009. Child health and living at high altitude. *Arch Dis Child* 94:806-811.
- Olaguer EP. 2012. The potential near-source ozone impacts of upstream oil and gas industry emissions. *J Air Waste Manag Assoc* 62:966-977.
- Parker SE, Mai CT, Canfield MA, Rickard R, Wang Y, Meyer RE, et al. 2010. Updated national birth prevalence estimates for selected birth defects in the United States, 2004–2006. *Birth Defects Res A Clin Mol Teratol* 88:1008-1016.
- Pétron G, Frost G, Miller BR, Hirsch AI, Montzka SA, Karion A, et al. 2012. Hydrocarbon emissions characterization in the Colorado front range: A pilot study. *J Geophys Res: Atmospheres* 117: D04304.
- Ren A, Qiu X, Jin L, Ma J, Li Z, Zhang L, et al. 2011. Association of selected persistent organic pollutants in the placenta with the risk of neural tube defects. *Proc Natl Acad Sci USA* 108:12770-12775.
- Ritz B, Yu F, Fruin S, Chapa G, Shaw GM, Harris JA. 2002. Ambient air pollution and risk of birth defects in Southern California. *Am J Epidemiol* 155:17-25.
- Ritz B, Wilhelm M, Hoggatt KJ, Ghosh JKC. 2007. Ambient air pollution and preterm birth in the environment and pregnancy outcomes study at the University of California, Los Angeles. *Am J of Epidemiol* 166:1045-1052.

- Ritz B, Wilhelm M. 2008. Ambient air pollution and adverse birth outcomes: Methodologic issues in an emerging field. *Basic Clinl Pharmacol Toxicol* 102:182-190.
- Romitti P, Sun L, Honein M, Reefhuis J, Correa A, Rasmussen S. 2007. Maternal periconceptual alcohol consumption and risk of orofacial clefts. *Am J Epidemiol* 166:775-785.
- Shepard T. 1995. Agents that cause birth defects. *Yonsei Med J* 36:393-396.
- U.S. EIA (United States Energy Information System). 2011a. International energy outlook 2011. DOE/EIA-0484(2011). [http://www.eia.gov/forecasts/ieo/pdf/0484\(2011\).pdf](http://www.eia.gov/forecasts/ieo/pdf/0484(2011).pdf). [accessed 22 May 2013].
- U. S. EIA. 2011b. Review of emerging resources: U.S. Shale gas and shale oil plays. <http://www.eia.gov/analysis/studies/usshalegas/pdf/usshaleplays.pdf>. [accessed 22 May 2013].
- Vidas H, Hugman B. 2008. Availability, economics, and production potential of North American unconventional natural gas supplies. Retrived from <http://www.ingaa.org/File.aspx?id=7878>. [accessed 22 May 2013].
- Vrijheid M, Martinez D, Manzanares S, Dadvand P, Schembari A, Rankin J, et al. 2011. Ambient air pollution and risk of congenital anomalies: A systematic review and meta-analysis. *Environ Health Perspect* 119 (5): 598-606.
- Wald N, Sneddon J. 1991. Prevention of neural tube defects: Results of the medical research council vitamin study. *Lancet* 338:131.
- Walther E. 2011. Screening health risk assessment Sublette County, Wyoming. SR2011-01-03. Pinedale, WY: Sierra Research, Inc. <http://www.sublettewyo.com/DocumentCenter/Home/View/438>. [accessed 22 May 2013].
- Wennborg H, Magnusson L, Bonde J, Olsen J. 2005. Congenital malformations related to maternal exposure to specific agents in biomedical research laboratories. *J Occup Environ Med* 47:11-19.
- Witter R, McKenzie L, Stinson K, Scott K, Newman L, Adgate J. 2013. The use of health impact assessment for a community undergoing natural gas development. *Am J Public Health* 103 (6): doi: 10.2105/AJPH.2012.301017.

Zielinska B, Fujita E, Campbell D. 2011. Monitoring of emissions from Barnett shale natural gas production facilities for population exposure assessment. Desert Research Institute.

<https://sph.uth.edu/mleland/attachments/Barnett%20Shale%20Study%20Final%20Report.pdf>

f. [accessed 22 May 2013].



**Table 1.** Study population characteristics for unexposed and exposed subjects from rural Colorado 1996-2009.

Maternal or Infant Characteristic	Total	Referent Group (No Wells within 10 miles)	Low (First Tertile) <sup>a</sup>	Medium (Second Tertile) <sup>a</sup>	High (Third Tertile) <sup>a</sup>
<b>Total n (%)</b>	124,842	66,626 (53)	19,214 (15)	19,209 (15)	19,793 (16)
Median	27	27	26	27	27
25 <sup>th</sup> Percentile	22	22	21	22	23
75 <sup>th</sup> Percentile	32	32	30	31	31
<b>Maternal Ethnicity (%)<sup>b</sup></b>					
White – Non-Hispanic	73	74	72	76	69
<b>Gender (%)</b>					
Male	51	51	51	51	51
<b>Maternal smoking (%)<sup>c</sup></b>					
Smokers	11	11	14	13	8
<b>Maternal Alcohol (%)</b>					
No	99	98	99	99	99
<b>Parity (%)</b>					
0	33	33	31	32	32
1	23	23	24	24	25
2	19	19	20	19	20
> 2	25	25	26	25	24
<b>Residential Elevation (feet)</b>					
Median	5000-5999	6000-6999	< 5000	5000-5999	< 5000
25 <sup>th</sup> Percentile	< 5000	5000-5999	< 5000	< 5000	< 5000
75 <sup>th</sup> Percentile	7000-7999	7000-7999	5000-5999	6000-6999	5000-5999
<b>Maternal Education (%)</b>					
< 12 years	21	20	26	19	22
12 years	30	30	33	29	28
13-15 years	23	22	25	25	24
≥ 16 years	26	28	18	26	27

<sup>a</sup>First tertile, 1 to 3.62 wells per mile; second tertile, 3.63 to 125 wells per mile; third tertile, 126 to 1400 wells per mile. <sup>b</sup>Maternal Ethnicity include White Non-Hispanic and

White Hispanic. <sup>c</sup>Smoking and alcohol during pregnancy

**Table 2.** Association between inverse distance weighted well count within ten-mile radius of maternal residence and congenital heart defects, neural tube defects, and oral clefts.

Inverse Distance Weighted Well Count <sup>a</sup>	0 Wells within 10 Miles	Low	Medium	High	Cochran-Armitage Trend Test p-value <sup>d</sup>
Live Births (N)	66,626	19,214	19,209	19,793	
<b>CHDs</b>					
Cases (N)	887	281	300	355	
Crude OR	1	1.1	1.2	1.3	< 0.0001
Adjusted OR (95% CI) <sup>b</sup>		1.1 (0.93, 1.3)	1.2 (1.0, 1.3)	1.3 (1.2, 1.5)	
<b>NTDs</b>					
Cases (N)	27	6	7	19	
Crude OR	1	0.77	0.90	2.4	0.01
Adjusted OR (95% CI) <sup>c</sup>		0.65 (0.25, 1.7)	0.80 (0.34, 1.9)	2.0 (1.0, 3.9)	
<b>Oral Clefts</b>					
Cases (N)	139	31	41	40	
Crude OR	1	0.77	1	0.97	0.9
Adjusted OR (95% CI) <sup>b</sup>		0.65 (0.43, 0.98)	0.89 (0.61, 1.3)	0.82 (0.55, 1.2)	

<sup>a</sup>Low = first tertile, 1 to 3.62 wells per mile, medium = second tertile, 3.63 to 125 wells per mile, high = third tertile, 126 to 1400 wells per mile. <sup>b</sup>Adjusted for maternal age, ethnicity, smoking, alcohol use, education, and elevation of residence, as well as infant parity and gender. <sup>c</sup>Adjusted only for residence elevation because of low numbers.

<sup>d</sup>Cochran-Armitage trend test performed as two tailed test on unadjusted logistic regression.

**Table 3.** Association between inverse distance weighted well count within ten-mile radius of maternal residence and preterm birth, term low birth weight.

<b>Inverse Distance Weighted Well Count<sup>a</sup></b>	<b>0 Wells within 10 Miles</b>	<b>Low</b>	<b>Medium</b>	<b>High</b>	<b>Cochran-Armitage Trend Test p-value<sup>d</sup></b>
<b>Preterm Birth</b>					
Live Births (N)	65,506	18,884	18,854	19,384	
Cases (N)	4,849	1,358	1,289	1,274	
Crude OR	1	0.97	0.92	0.88	< 0.0001
Adjusted OR (95% CI) <sup>b</sup>		0.96 (0.89, 1.0)	0.93 (0.87, 1.0)	0.91 (0.85, 0.98)	
<b>Term Low Birth Weight</b>					
Full Term Live Births (N)	60,653	17,525	17,565	18,104	
Cases (N)	2,287	525	471	432	
Crude OR	1	0.79	0.70	0.62	< 0.0001
Adjusted OR (95% CI) <sup>b</sup>		1.0 (0.9, 1.1)	0.86 (0.77, 0.95)	0.9 (0.8, 1)	
<b>Mean difference in birth weight (grams)<sup>c</sup></b>	0	5 (-2.2, 13)	24 (17, 31)	22 (15, 29)	

<sup>a</sup>Low = first tertile, 1 to 3.62 wells per mile, medium = second tertile, 3.63 to 125 wells per mile, high = third tertile, 126 to 1400 wells per mile <sup>b</sup>Adjusted for maternal age, ethnicity, smoking, alcohol use, education, and elevation of residence, as well as infant parity and gender. <sup>c</sup>Adjusted for maternal age, ethnicity, smoking, alcohol use, education, and elevation of residence, as well as infant parity, gender, and gestational age. <sup>d</sup>Cochran-Armitage trend test performed as two tailed test on unadjusted logistic regression.

**Table 4.** Association between inverse distance weighted well count within ten-mile radius of maternal residence and congenital heart defect diagnostic groups.

Inverse Distance Weighted Well Count <sup>a</sup>	0 Wells within 10 Miles	Low	Medium	High
<b>Conotruncal Defects</b>				
Cases (N)	40	14	13	15
Adjusted OR (95% CI) <sup>e</sup>	1	1.1 (0.57, 2.2)	1.1 (0.55, 2.0)	1.2 (0.6, 2.2)
<b>Ventricular Septal Defects</b>				
Cases (N)	210	68	59	84
Adjusted OR (95% CI) <sup>b</sup>	1	1.3 (0.96, 1.8)	1.1 (0.81, 1.5)	1.5 (1.1, 2.1)
<b>Endocardial Cushion and Mitrovalve Defects</b>				
Cases (N)	39	14	12	12
Adjusted OR (95% CI) <sup>e</sup>		0.81 (0.42, 1.6)	0.80 (0.41, 1.5)	0.67 (0.33, 1.32)
<b>Pulmonary Artery and Valve Defects</b>				
Cases (N)	137	52	62	66
Adjusted OR (95% CI) <sup>b</sup>	1	1.3 (0.89, 1.8)	1.5 (1.1, 2.1)	1.6 (1.1, 2.2)
<b>Tricuspid Valve Defects</b>				
Cases (N)	9	5	8	8
Adjusted OR (95% CI) <sup>e</sup>	1	2.6 (0.75, 9.1)	3.9 (1.3, 11)	4.2 (1.3, 13)
<b>Aortic Artery and Valve Defects</b>				
Cases (N)	75	22	21	24
Adjusted OR (95% CI) <sup>b</sup>	1	1.1 (0.68, 1.9)	1.0 (0.62, 1.8)	1.2 (0.73, 2.1)
<b>Patent Ductus Arteriosus</b>				
Cases (N)	59	18	17	15
Adjusted OR (95% CI) <sup>b</sup>	1	1.0 (0.56, 1.8)	0.96 (0.55, 1.7)	0.83 (0.44, 1.5)

<sup>a</sup>Low = first tertile, 1 to 3.62 wells per mile, medium = second tertile, 3.63 to 125 wells per mile, high = third tertile, 126 to 1400 wells per mile. <sup>b</sup>Adjusted for maternal age, ethnicity, smoking, alcohol use, education, and elevation of residence, as well as infant parity and gender. <sup>c</sup>Adjusted only for residence elevation of because of low numbers.